

Towards a Simple Gaussian Model to Describe Multiple Source Areas during Wind Erosion Events.

H.J. Butler: Department of Mathematics & Computing, U.S.Q. Toowoomba, Queensland, 4350.

W.L. Hogarth: Faculty of Environmental Science, Griffith University, Nathan, Queensland, 4111.

G.H. McTainsh: Faculty of Environmental Science, Griffith University, Nathan, Queensland, 4111.

Abstract Wind erosion is a broad-scale process in inland Australia. When conditions are conducive to wind erosion, dust storms can entrain fine sediment over large areas of the continent, however closer examination indicates that the dust source areas are often spatially discrete. The fine sediment entrained from these sources, is transported as dust plumes, which may coalesce at some point downwind. While some progress has been made in estimating the dust load in these plumes [Knight et al. 1995], the accuracy of these estimates is limited by the size and shape of the source region assumed. In addition, soil loss per unit area is a more appropriate measure of soil erosion than total plume load, but estimating loss per unit area requires accurate estimation of source areas. The new model developed hopes to overcome these limitations by working from the source area downwind, rather than back-tracking to estimate the source area as done by Knight et al. [1995]. As a result of this, the new model is quite distinct from that of Knight et al. [1995], in that it no longer assumes that entrainment is uniform across a single source area, and consequently that the concentration profile is uniform across the source area. The new model uses a Gaussian plume model [Zannetti 1990], with the dispersion parameters based on the Hanna et al. [1982] estimates for rural areas. As a first approximation, the model describes dust loads emanating from sources of different strength and spacing along a crosswind line. This configuration is indicative of spatially discrete sources, with different soil erodibility and cover protection, producing different source strengths. Initial results from the model indicate that the nature of the downwind dust concentration profile is dependent on the following factors: downwind distance from source, source strengths, and crosswind source separation. A detailed discussion of the relationship between the above factors is presented. As a result of this discussion a number of conjectures are made about the nature of the physical system.

1. INTRODUCTION AND BACKGROUND

Wind erosion is a major geomorphic process in much of inland Australia, especially when drought conditions prevail and adequate ground cover levels are hard to maintain. In any given wind erosion event, much of the fine material picked up by the wind is transported from the source in dust plumes [McTainsh 1989]. Since the amount of material available to the wind for entrainment is limited by factors such as soil erodibility, ground cover and soil moisture, the resultant dust plume (cloud) could be made up of material from a number of discrete sources. The number, location and size of these source areas has, to date, proved difficult to describe.

While some progress has been made in modelling dust plumes in Australia [Knight 1990, Knight et al. 1995] and overseas [D'Almeida 1986, McMahon et al. 1976, Foda 1983]. The majority of these models lack the ability to describe spatial variability in dust concentration in the plume and use backtracking techniques to describe the source areas (see figure 1).

The Composite Box Model of Knight et al. [1995], is the only model to be currently applied to Australian conditions. In the originating source box (Box 1) (i.e. the Birdsville Box see figure 2) the whole source area covered by the box is assumed to be entraining at constant rate. The only other source box assumed to be entraining dust is the Charleville box (Box 3) and here entrainment is limited to only a certain percentage of the area of the box. The amount of material entrained is estimated by using the amount produced if the total area was entraining, then taking a percentage of this. It is further assumed that the dust concentration within each box is constant.

While this last assumption "maybe" consistent with the dust concentration profile observed in the Brisbane box (Box 4), where there is little variation in dust concentration across the box, its validity in and around the Charleville box must be questioned. In the Charleville box, we have a situation where only discrete regions of the box are eroding and producing dust plumes. Around each of these regions one would expect a substantial increase in the concentration of dust in the air (i.e. local hot spots). Clearly then this assumption is in-

adequate in its description of dust concentrations within and around boxes that contain discrete source regions.

In order to describe spatial variations of dust concentrations around individual source regions, it is necessary to describe the dust concentrations that may result from discrete sources that may occur in that region. To achieve this the source model must be able to treat each source as a discrete identity, and predict how these sources will intermix with the given

source region.

The use of backtracking techniques in current models provides a means of determining the boundaries of the source regions, it however doesn't provide an estimate of how the sources are scattered within the region. Determination of how much of this region is "eroding", is left to the researchers to estimate based on their knowledge of the regions environmental conditions.



Figure 1: Calculated back-trajectories of the 1987 event that effected South-East Queensland, Australia (after Knight et al. [1995]).




Figure 2: The location of the various Boxes used by Knight et al. [1995].

This estimate has to include at least two factors:-

- the strength of each of the local source areas in the source region during the event.
- the percentage area of the source region that is entraining during an event.

Reasonably accurate estimates of these factors can be made if field data is collected for an event [McTainsh et. al. 1995], but if this is not available a fair amount of guess work, based on passed experience of the researcher is used to provide estimates for these factors on an event basis. Our source model in final form will allow the strength of each source to be defined separately from calculations based upon measured environmental conditions. Field data will also allow better estimates of the location of local source areas on an event basis to be made. This data will also be used, along with sensitivity analysis to determine a suitable source profile (source distribution) for the model that best describes the reality of the situation.

2. THE SOURCE MODEL

The source model is based on a Gaussian Plume Dispersion model (see Zannetti [1990]), with the following assumptions:

- local source areas have a constant source strength with time $Q(\mu\text{gs}^{-1})$.
- The average dust concentrations in the vertical and crosswind directions follow a Gaussian distribution.
- The effective height of the source above the ground is 0 m (i.e sources are ground based and there is no thermal or turbulent plume rise away from the surface).
- Wind speed (u) is uniform throughout the source area.
- No deposition occurs in the source region (this will be relaxed following future model development).
- All material entrained is reflected from the surface i.e there is no absorption of entrained material by the surface (this will be relaxed once experimental data becomes available).

Given these assumptions, Hanna et al. [1982] state that the concentration $C(\mu\text{gm}^{-3})$ at any point downwind of a point source is given by:-

$$C(x, y, z) = \frac{Q}{\pi\sigma_y\sigma_z u} \exp\left(-\frac{y^2}{2\sigma_y^2}\right) \exp\left(-\frac{z^2}{2\sigma_z^2}\right) \quad (1)$$

where z is the height above the ground, y is the cross wind position from some reference point and $\sigma_y(x)$ and $\sigma_z(x)$ are the standard deviations (Dispersion Parameters) of the concentration in the y and z directions respectively.

In choosing the functional form of the dispersion parameters ($\sigma_y(x)$ and $\sigma_z(x)$), it is assumed that they are functions of only the downwind distance x and the stability of the system.

The stability of the system is included in the model by using pasquill-stability categories outlined by Hanna et al. [1982], as a basis for classifying the turbulent stability of the system. Once the stability category for the system has been determined, then the functional form of the dispersion parameters (which are only functions of downwind distance) is chosen for the given stability conditions.

As an initial approximation, it is assumed that they have a similar functional form to that outlined by Zannetti [1990] for neutral rural meteorological conditions. Mathematically they take the following form:-

$$\sigma_y(x) = \frac{0.08x}{\sqrt{1 + 0.0001x}} \quad (2)$$

and

$$\sigma_z(x) = \frac{0.06x}{\sqrt{1 + 0.0015x}} \quad (3)$$

As an initial approximation, local area sources are assumed to be line sources located along a cross-wind line. Each distinct line source is assumed to consist of a number of distinct point sources. This is done for a variety of reasons, such as:-

1. it allows the functional form of the source strength across any source region to be easily altered (i.e it is possible to have a line source with a higher strength in the centre to that observed at the ends.) This situation is more representative of natural occurring sources.
2. If at any stage in the future it is required that sources be angled to the wind, then this can be implemented in the model, without substantial modifications.

By taking this approach, each local source area can be assigned a unique source strength (currently this is still assumed to be constant across each source in our model) and separation distance from its neighbour.

2.1 The Functional Forms of $\sigma_y(x)$ and $\sigma_z(x)$

While the functional forms of $\sigma_y(x)$ and $\sigma_z(x)$, represent a reasonable starting point for the purpose of developing and testing the model, their relevance to Australian conditions, must at some stage be consolidated, in terms of some of the concepts outlined below.

Since the purpose of this model is to describe a dispersion process that involves multi-sized particles of differing specific densities, the very assumption that the dispersion parameters are independent of the physical properties of the particles, needs to be questioned. In order to change this assumption, the turbulent forces acting on the various particles must be sufficiently different, to have a noticeable effect on the dispersion of the particles. However, at present there is insufficient experimental data on particle size related dispersion effects, from which to base any change.

It is also obvious that the actual functional form of the dispersion parameters may also need to be revised along the following lines:-

- Do Australian soil particles behave similarly to those used in US based experiments for determining the dispersion parameters?

- Are atmospheric dispersion conditions in inland Australia similar to those in the US (i.e. do we have more temperature-induced turbulent mixing, compared with the US)?

Once these issues are resolved with empirical data the functional forms of the dispersion parameters can be changed to better describe Australian conditions.

3. THE PROBLEM OF SCALES

The broad-scale nature of wind erosion and dust transport, means that we should describe the process at three distinct spatial scales (i.e. the local, regional and continental scale). Eventually it may be possible to describe these in one model, rather than using separate models to describe essentially the same process. The major problem with using the same approach in each case, is picking how to define the source area at each scale.

To understand why this is such a major problem consider a source area made up of numerous discrete sources (as in figure 3). If an observer is located at a sufficient distance downwind of the source, a number of the dust plumes would have coalesced and appear as single concentration profile. Thus if you are trying to describe the concentration at some point after this has occurred a single source initially could be used to describe the sources that coalesced. The point where coalescing occurs is dependent on three factors: the dispersion parameters, source separation and the relative strengths of each source. Therefore picking adequate source descriptions for each scale will depend on our understanding of how these factors interact. Much of this understanding will come through the work, being done on our current model.

4. RESULTS FROM THE MODEL

4.1 Separation of Sources

Source separation has a direct effect on the downwind distance before the plumes from each source coalesce into a single profile. During transition from multiple plumes to a single plume the model predicts that there will be three distinct transitional stages (see Figure 4). These stages are outlined below:-

1. Each source has a distinct peak in the profile, that is clearly associated with that source.
2. A plateau (or shoulder, see discussion in section 4.2) is apparent in the profile.
3. The profile is similar to that generated from a single point source.

Between each of these distinct profiles there is a transition profile.

Each of these stages represents various physical stages in the model. The first stage, where each source has a distinct peak, is representative of the where the source plumes mix only slightly. This mixing gradually increases with downwind distance until the centre of the plumes are beginning to mix, so that a plateau is produced around the two centres (if sources are of equal strength, otherwise a shoulder is produced see figure 4 and figure 5.) The mixing gradually increases with downwind distance until the plumes have totally combined and are indistinguishable from a single point source "profile" (i.e. a simple gaussian distribution).

4.2 Source Strength Effects

While the separation of the sources appears to be the major controlling influence on the distance downwind when the plumes coalesce, source strength also plays a role. If one source is significantly stronger than the other, the weaker has little effect on the major source, except to produce a distinct tail in the profile, depending on separation of the sources. This distortion in the profile soon becomes insignificant as the sources mix further, especially when compared to the concentration change in the region of the stronger source (i.e. the effect is still present but produces only a small change in concentration when compared to changes due to the dispersion of material near the stronger source).

If however, the sources are of similar (but not equal) strength the stages outlined in 4.1 are quite distinct (see Figure 5). Here the shoulder is produced by the mixing of two different strength sources.

Source strength also has an effect on the plateau stage. As mentioned earlier, if the same separation is maintained, but with different source strengths the plateau effect reverts to a shoulder/step effect. However if the source strength remains equal but is increased (or decreased), the profile produced represents an increase (or decrease) in the concentration present across the profile, but the plateau effect is maintained (see figure 6).

In the model while the source strength is increased, the cross-wind dispersion remains constant. Thus the two plumes coalesce at the same point, producing the plateau. However whether this phenomena actually occurs in the physical system must be questioned. The reason this may not be the case, is that the more material present in the plume, the higher the inter-particle forces and the more collisions that will occur in the plume. Then if sufficient material is

present in the cloud, the increase in these two factors should induce greater dispersion rates within the plume. The opposing argument is that inside the plume restoration forces will balance out any increase in these two factors.

4.3 Downwind Effects

As you would expect from the above, both the separation and the strength play an important role in the effects seen in the downwind profiles. These variables control not only when the effect is seen, but also the strength of the effect.

The nature of the effect is seen in figure 7, which is a downwind profile taken half way between two line sources. As can be seen clearly in this profile there is a gradual build up in the concentration with downwind distance until a maximum concentration is reached, and then there is a gradual decrease




Figure 3: A pictorial representation of a possible distribution of area sources within two distinct source regions.




Figure 5: Crosswind profiles illustrating the effects of source strength on the resultant crosswind profile. Source Details: two 1km line sources, with the source strength of second being 0.75 the first. Profile taken at: 10km downwind of the source line and at a height of 10m.

as the plumes continue to disperse and mix. This is in contrast to downwind profiles taken in the centre of one of the sources, which are illustrated in figure 8. Here all that is evident as the two sources combine, is a kink in the profile.

This behaviour is consistent with what one would expect as the two sources combine. In the centre of two sources one would expect an initial buildup in material, as there is initial no material in this region and as material is transported into the region the concentration should increase. This increase should continue until the two sources are well mixed, after this point is reached the material is still dispersing, so there should be a gradual decrease in the concentration as the model predicts. While in the centre of a source, the effect of the other source beginning to mix should be a slight increase in the material present in the plume at that point, thus causing the concentration to slow its rate of decrease (or to actually start increasing) at that point.




Figure 4: Illustration showing the effects of three different source separations on the concentration profile. Source Details: two 1km line sources of equal strength. Profile taken at: 10km downwind of the source line and at a height of 10m.




Figure 6: illustration of the effect of increasing the source strength of both sources equally on the plateau produced when both sources are equal. Source Details: two 1km line sources of equal strength. Profile taken at: 10km downwind of the source line and at a height of 10m.




Figure 7: The downwind concentration profile taken halfway between the two sources. Source Details: two 1km line sources of equal strength. Profile taken at: a height of 10m.




Figure 8: The downwind concentration profile taken in the centre of the first source. Source Details: two 1km line sources of equal strength. Profile taken at: a height of 10m.

5. IMPLICATIONS OF THESE RESULTS

While the source model we have outlined here is crude in many respects, it does however provide some insight into the behaviour of dust plumes from multiple local dust source areas. It is obvious from the initial results of the model that care must be taken in the description of sources if accurate results are to be achieved. The eventual positioning of local source areas will of course be determined by the field data for the source region. The model also indicates that source properties (separation etc.) must be included in any transport model if accurate regions of peak concentrations are to be predicted from the model.

In sections 2 and 4 a number of assumptions are raised concerning aspects of the model. These areas of the model require significantly more research before they can be relaxed. However, they do provide a reasonable initial framework for the model.

6. WHERE TO FROM HERE ?

Over the next six months, experimental field and laboratory data will be collected to verify aspects of the model in its current form and model changes will be made based on these results. Also during this time it is planned to introduce dry and wet deposition into the model, to account for these two processes over the source area.

Once the source model is finalised, it is planned to use it as a basis for a long range dust transport model. This new model will incorporate multiple source areas and spatial variations in concentrations. If successful this model will for the first time allow spatial predictions of dust concentrations to be

made on the East Coast of Australia, for major dust events. This will mean forecasters will be able to make predictions of areas where the visibility will be most effected during major dust events, thus allowing more precise local raised dust warnings to be issued.

7. SUMMARY

The Gaussian model outlined here, provides a useful method to model spatial concentrations of dust within a given source region. The model does however have inherent problems in terms of a number of unrealistic assumptions. However, until better experimental data from current studies become available, it is extremely hard to improve on these assumptions.

In particular, the current model also lacks one vital component in any transportation model; a deposition term. However, as this component of the model controls or influences much of the physical behaviour of the model, particular care must be taken in its inclusion model. Current experimental work on deposition is centred around obtaining accurate theoretical/empirical estimates of deposition velocity and washout rates of dust particles. Once this work is completed deposition will form an integral part of the source model.

8. ACKNOWLEDGEMENTS

The authors acknowledge the support of the Australian Research Council in funding both the theoretical and experimental aspects of this research.

9. TERMINOLOGY

backtracking A technique used in meteorology to back-track the path of air parcels over a given period.

source region A area of land that contains a number of discrete sources (see figure 3).

source area A term used to denote the smallest discrete source (see figure 3)

separation of sources The distance between line sources which is taken to be the distance from the trailing edge of one source to beginning of the next source.

entrainment The term given to the process of particles being ejected into the air.

10. REFERENCES

D'Almeida, G. A., A model for Saharan dust transport, *Journal of Climate and Applied Meteorology*, 25(7), 903-916, 1986.

Foda, M. A., Dry-fall of fine dust on sea, *Journal of Geophysical Research*, 88, 6021-6026, 1983.

Hanna, S., G. Briggs, and R. Hosker, *Handbook on Atmospheric Diffusion*, Technical Information Center U.S Department of Energy, 1982.

Knight, A., Estimates of long distance dust transport over Eastern Australia, Honours Dissertation, Griffith University, 1990.

Knight, A., G. McTainsh, and R. Simpson, Sediment loads in an Australian dust storm: Implications for present and past dust processes. *CATENA (in press)*, 1995.

McMahon, T. A., P. Denison, and R. Fleming, A long-distance air pollution transport model incorporating washout and dry deposition components, *Atmospheric Environment*, 10, 751-761, 1976.

McTainsh, G., Quaternary aelian dust processes and sediments in the Australian region, *Quaternary Science Reviews*, 8, 235-253, 1989.

McTainsh, G., W. Nickling, J. Leys, and A. Lynch, Wind erosion rates and processes from field measurements in the channel country, Queensland. In preparation, 1995.

Zannetti, P., *Air Pollution Modelling: Theories, Computational Methods and Available Software*, Van Nostrand Reinhold, 1990.