

Importance of Dispersion for Shoaling Waves

J. Pitt^a, C. Zoppou^a, S.G Roberts^a

^a*Mathematical Sciences Institute, Australian National University, Canberra, ACT 0200, Australia*
Email: jordan.pitt@anu.edu.au

Abstract: A tsunami has four main stages of its evolution; in the first stage the tsunami is generated, most commonly by seismic activity near subduction zones. The second stage is the tsunamis propagation through the ocean far from the coast, where variation in bathymetry is slight and gradual. The third stage is the shoaling and interaction of the tsunami with bathymetry as it approaches the coastline. Finally the tsunami reaches and inundates the shore. For our purposes the hydrodynamic models we are interested in deal with the final three stages of the evolution of a tsunami.

The propagation of a tsunami with wavelength λ through water that is H deep is well understood when $\lambda/H \leq 1/20$, which we call shallow water as noted by Sorensen (2006). The wavelengths for tsunamis range from a few to hundreds of kilometres, while the maximum water depth is $11km$ at the Marianas trench, so that most tsunamis occur in shallow water. This stage of tsunami behaviour is adequately modelled using the shallow water wave equations. Current research into tsunamis focuses around more complex approximations to the Euler equations for the third and fourth stages. In this paper we focused on the Serre equations as they are considered a very good model for fluid behaviour up to the shoreline, and they reduce to the shallow water wave equations for large wavelengths.

Although more complicated, the Serre equations provide a better description of the fluid behaviour than the shallow water wave equations and are therefore more computationally expensive to solve numerically. In particular for the methods of this work, we find that the Serre equations have a run-time 50% longer than our equivalent finite volume method for the shallow water wave equations in the one dimensional case. To simulate tsunamis as efficiently as possible it is important to know when using the more complicated Serre equations leads to more accurate predictions of the evolution of a tsunami than the shallow water wave equations. To investigate this we have numerically simulated a laboratory experiment of periodic waves propagating over a submerged bar, and the propagation of a small amplitude wave up a gradual linear slope using both the Serre and the shallow water wave equations.

The results of these simulations demonstrated that the Serre and shallow water wave equations produce similar results for shoaling waves when the wavelength is large compared to the water depth. This is not surprising as this is the regime under which the shallow water wave equations are derived. However, outside this regime the shallow water wave equations are a poor model for wave shoaling and propagation, poorly approximating the shape and maximum height of waves. Furthermore we demonstrate that for steep waves generated by shoaling, the shallow water wave equations can underestimate the arrival time and amplitude of an incoming wave. These results suggest that for a tsunami it is sufficient to use the shallow water wave equations in stages two and some of stage three, even for large changes in bathymetry. Although dispersive equations such as the Serre equations are required to accurately capture fluid behaviour in stages three and four nearer to the coastline, particularly when wavelengths are short or waves are steep. Since the Serre equations represent only a moderate increase in run-times this suggests that our inundation models should be based on them.

Keywords: *Tsunamis, shoaling waves, dispersion, models, shallow water wave equations*

1 INTRODUCTION

The interaction of waves with bathymetry and particularly the shoaling of waves plays a central role in modelling the inundation caused by a tsunami. Current models for tsunami inundation such as ANUGA utilise the non-dispersive shallow water wave equations. Current research into tsunami inundation includes building numerical models based on dispersive equations, such as the Serre equations. The Serre equations allow for more detailed fluid behaviour such as dispersion, which is the process by which waves of different frequencies travel at different speeds. The Serre equations are considered by Bonneton *et al.* (2011) to be the most appropriate approximation to the Euler equations near the shoreline. The one-dimensional Serre equations for a depth of water $h(x, t)$ over a bed profile defined by $b(x)$ travelling at velocity $u(x, t)$ are given by Zoppou *et al.* (2017) as

$$\frac{\partial h}{\partial t} + \frac{\partial(uh)}{\partial x} = 0 \quad (1a)$$

and

$$\frac{\partial uh}{\partial t} + \frac{\partial}{\partial x} \left(u^2 h + \frac{gh^2}{2} + \frac{h^3}{3} \Phi + \frac{h^2}{2} \Psi \right) + \left(gh + \frac{h^2}{2} \Phi + h\Psi \right) \frac{\partial b}{\partial x} = 0 \quad (1b)$$

where

$$\Phi = \left(\frac{\partial u}{\partial x} \right)^2 - u \frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial x \partial t}, \quad \Psi = \frac{\partial u}{\partial t} \frac{\partial b}{\partial x} + u \frac{\partial u}{\partial x} \frac{\partial b}{\partial x} + u^2 \frac{\partial^2 b}{\partial x^2}.$$

Setting $\Phi = \Psi = 0$, corresponding to long wavelengths transforms the Serre equations into the shallow water wave equations. The additional Φ and Ψ terms make the Serre equations more complex and therefore more computationally difficult to solve. However, the additional behaviour they allow for is not always significant, in particular when wavelengths λ are long compared to the water depth H such that $H/\lambda \leq 1/20$. In this paper we will investigate the impact of these extra terms by comparing the numerical solution of the Serre equations with the solution of the shallow water wave equations. To do this we will use a well-validated numerical method for the Serre equations by Zoppou *et al.* (2017) and the ANUGA software for the shallow water wave equations. We begin by simulating an experiment of Beji and Battjes (1994) where periodic waves travelled over a submerged bar, and then move to a more appropriate scenario for tsunamis, the propagation of a long wavelength, small amplitude solitary wave up a uniformly sloping beach.

2 PERIODIC WAVE OVER A SUBMERGED BAR

In the series of experiments undertaken by Beji and Battjes (1994) periodic waves travel over a submerged trapezoidal bar. Their wave tank was 37.7m in length, 0.8m wide with walls 0.75m high with a constant water surface sitting at 0.4m. A diagram of this wave tank displaying the bed profile, water surface and locations of wave gauges is presented in Figure 1. Waves were generated by a wave maker located at $x = 0$ m, and were absorbed downstream by a gravel beach that extends to the edge of the wave tank. There were seven wave gauges that recorded the depth of water over time at the following locations; WG1: 5.7m, WG2: 10.5m, WG3: 12.5m, WG4: 13.5m, WG5: 14.5m, WG6: 15.7m and WG7: 17.3m.

In this paper we focus on the propagation of periodic non-breaking sinusoidal waves with high frequency over the bar. These waves had a period of $T = 1.25$ s, a wavelength of $\lambda = 2.05$ m and an amplitude of $A = 0.0125$ m. These waves travelled without deformation towards the bar. The waves steepened as they travelled up the front slope of the bar due to shoaling. On the horizontal top of the bar higher frequency waves are slowly released. The release of these waves is accelerated as the waves travel leeward.

The numerical results for the Serre equations and the shallow water wave equations are presented in Figure 2. We present only wave gauges 2 through 4, as the deficiencies of the shallow water wave equations in simulating this experiment are already apparent by the 4th wave gauge.

In these experiments the time series data from wave gauge 1 for the water depth is used as the left boundary condition to simulate the incoming wave. Both models reproduce it precisely and we include it here as a reference for the incoming wave profile. It can be seen that the Serre equations reproduce the experiment results well. The shallow water wave equations results are poor for both the shape and speed of the waves.




Figure 1. Wave tank diagram with wave gauges marked by number showing the water surface (—) and the bed profile (—) .

This is because the shallow water wave equations are appropriate when $H/\lambda \leq 1/20$, which is not the case in these experiments as $0.4m/2.05m \approx 1/5$. Therefore an increase in the wavelength of the experiment should result in a better reproduction of the results by the shallow water wave equations.

In another experiment of Beji and Battjes (1994) the propagation of longer wavelength sinusoidal waves with a period of $T = 1.25s$, a wavelength of $\lambda = 3.69m$ and an amplitude of $A = 0.01m$ over the same submerged bar was observed. The results for using the Serre equations and the shallow water wave equations to model this experiment are displayed in Figure 3. The shallow water wave equations model these waves better than the waves of the previous experiment because they are longer. Unfortunately $0.4m/3.69m \approx 1/9$, so these waves were still too short to be appropriate for the shallow water wave equations, and so the results are inaccurate.

This experiment is specifically designed to study dispersive effects, and so the failure of the shallow water wave equations is not surprising. However, the inaccuracy of the shallow water wave equations was larger than anticipated. This demonstrates that dispersion can be a significant factor in the shoaling of waves, with large differences between the simulations using the Serre and shallow water wave equations.

3 PROPAGATION OF A SOLITON OVER A SLOPE.

The numerical results for the propagation of periodic waves over a submerged bar demonstrated large differences between the Serre and shallow water wave equations. These differences are pronounced due to the fact that the experimental set-up had waves with a wavelength that is smaller than 20 times the water depth, which is not appropriate for the shallow water wave equations. To rectify this another numerical experiment was performed simulating the propagation of a long wavelength, small amplitude wave over a gradual uniformly sloped bed. This scenario is analogous to a tsunami propagating from a continental shelf to the shoreline.

In this simulation the initial conditions are the soliton of the Serre equations from El et al. (2006) with an amplitude of $0.01m$ on top of quiescent water $1m$ deep. A soliton of the Serre equations was used because it has an appropriate wavelength for the shallow water wave equation and it is a solitary wave like a tsunami. The water depth is kept at a fixed $1m$ throughout the scenario. The bed profile is defined in the following way; $(x, b) = [(-100m, 0m), (100m, 0m), (149.5m, 0.99m), (250m, 0.99m)]$. The initial conditions and the numerical solutions for both the Serre equations and the shallow water wave equations are plotted in Figure 4.

It can be seen that the behaviour of the numerical solutions of the Serre equations and the shallow water wave equations are very similar when the wavelength is large compared to the depth. There is very little wave deformation for both equations after $t = 20s$. For the Serre equations the soliton travels without deformation due to a balance between non-linearity and dispersion. For the shallow water wave equations the soliton travels with little deformation because the effect of non-linearity is small, as the wave amplitude is small. This demonstrates the negligible effect of dispersion for long wavelength and small amplitude waves far from the coast.




Figure 2. Wave height readings from various wave gauges from the experiment of Beji and Battjes (1994) (—) compared to a simulation of the experiment using the Serre equations (\circ) and shallow water wave equations ($+$).




Figure 3. Wave height readings from various wave gauges from the experiment of Beji and Battjes (1994) (—) compared to a simulation of the experiment using the Serre equations (\circ) and shallow water wave equations ($+$).




Figure 4. The water profile from the numerical solution for the Serre equations (—) and the shallow water wave equations (—) for the soliton travelling over a slope, at different times. The initial conditions for both numerical models are also presented in terms of stage (—) and bed profile (—).

As the soliton progresses up the slope there is also very little difference between the numerical solutions throughout the shoaling process as can be seen at $t = 40s$. It is only when the waves become steep after the slope ends that differences begin to arise although both numerical solutions are quite similar at $t = 60s$ and at $100s$. This demonstrates that dispersive effects are negligible for shoaling in this case, but play a role for the steep waves generated by shoaling. The main difference between the Serre equations and the shallow water wave equations is the emergence of undulations trailing the bore that has formed. These undulations have been observed experimentally by Chanson (2009) and numerically by Mitsotakis et al. (2014) for the Serre equations and even for the Euler equations by Mitsotakis et al. (2017). These undulations oscillate around the bore of the shallow water wave equations, significantly increasing its maximum amplitude. These undulations cause an undular bore to travel slightly quicker than the bores of shallow water wave equations as demonstrated by Pitt et al. (2017). These results suggest that the shallow water wave equations will slightly underestimate the arrival time as well as the maximum amplitude of a steep advancing wave.

4 CONCLUSIONS

Well-validated numerical methods for the Serre and shallow water wave equations were used to investigate the effect of dispersion on shoaling waves. It was demonstrated that when the water is shallow, dispersion is negligible during shoaling. Conversely it was demonstrated that when the water is not shallow dispersion play a significant role in the waves interaction with bathymetry. It was also demonstrated that dispersion plays a significant role when steep gradients in the water surface are present, increasing the maximum amplitude and the speed of an incoming wave, compared to the non-dispersive equations. For tsunamis this is impor-

tant as shoaling waves steepen as they approach the coast line. These results demonstrate the need for the Serre equations to accurately capture fluid behaviour in the near shore and the adequacy of the shallow water wave equations for modelling long waves. Due to the Serre equations being only a moderate increase in computational expense we believe inundation models should be based on them.

REFERENCES

- Beji, S. and J. Battjes (1994). Numerical simulation of nonlinear wave propagation over a bar. *Coastal Engineering* 23(1), 1–16.
- Bonneton, P., E. Barthélemy, F. Chazel, R. Cienfuegos, D. Lannes, F. Marche, and M. Tissier (2011). Recent advances in Serre-Green Naghdi modelling for wave transformation, breaking and runup processes. *European Journal of Mechanics B/Fluids* 30(6), 589–597.
- Chanson, H. (2009). An experimental study of tidal bore propagation: The impact of bridge piers and channel constriction. Research Report CH74/09, School of Civil Engineering, The University of Queensland. 104p.
- El, G., R. H. J. Grimshaw, and N. F. Smyth (2006). Unsteady undular bores in fully nonlinear shallow-water theory. *Physics of Fluids* 18(2), 027104.
- Mitsotakis, D., D. Dutykh, and J. Carter (2017). On the nonlinear dynamics of the traveling-wave solutions of the serre system. *Wave Motion* 70(1), 166–182.
- Mitsotakis, D., B. Ilan, and D. Dutykh (2014). On the galerkin/finite-element method for the serre equations. *Journal of Scientific Computing* 61(1), 166–195.
- Pitt, J., C. Zoppou, and S. Roberts (2017). Behaviour of the serre equations in the presence of steep gradients revisited. *arXiv preprint arXiv:1706.08637*.
- Sorensen, M. R. (2006). *Basic Coastal Engineering* (3 ed.).
- Zoppou, C., J. Pitt, and S. Roberts (2017). Numerical solution of the fully non-linear weakly dispersive serre equations for steep gradient flows. *Applied Mathematical Modelling* 48, 70–95.